

Instruction sheet

AFFINIMIP® SPE Metanephrines cartridges

Format: 1mL

PURIFICATION PROCEDURE OF Metanephrines in plasma or serum

Users should read all instructions before using this kit.

For research use only (RUO)

AFFINIMIP® SPE Metanephrines is:

developed and manufactured by AFFINISEP

E-mail: contact@affinisep.com

Technical support: tech.support@affinisep.com

www.polyintell.com

Version 1.2



Table of contents

	PRII	NCIPLE OF AFFINIMIP®SPE	3
2	PRC	DDUCT INFORMATION	3
3	PRE	CAUTIONS FOR USE	3
4	GEN	NERAL INSTRUCTIONS FOR SPE PURIFICATION	3
	4.1	Materials and equipment required	3
	4.2	Flow rate	4
	4.3	Drying step	4
	4.4	Template use	4
5	PUR	IFICATION PROCEDURE OF METANEPHRINES FROM PLASMA OR SERUM	4
	5.1	Preparation of solutions	4
	5.2	Preparation of the loading solution from plasma or serum (dilution by 5)	4
	5.3	Protocol for purification of Metanephrines	5



Method for Selective Phase Extraction of Metanephrines using Molecularly Imprinted Polymers

1 PRINCIPLE OF AFFINIMIP® SPE

AFFINIMIP® SPE is a solid phase obtained by a polymerisation process to create a three-dimensional network that recognizes the shape and functional group positions of a template molecule. The **AFFINIMIP® SPE** selectivity comes from the technology of molecularly imprinted polymer (MIP) used during the synthesis.

AFFINIMIP® SPE Metanephrines has been developed to selectively extract Metanephrines from serum or plasma.

By using AFFINIMIP® SPE, the expected result is a clean-up and a pre-concentration of the sample at trace levels.

2 PRODUCT INFORMATION

Description of the kit

Each solid phase extraction (SPE) cartridge **AFFINIMIP® SPE Metanephrines** contains 30mg of sorbent in a 1mL cartridge.

Information and storage

Storage: Room temperature.

Each cartridge has a single use.

3 PRECAUTIONS FOR USE

SPE methods developed for C18 or other sorbents are not appropriated for AFFINIMIP® SPE Metanephrines.

4 GENERAL INSTRUCTIONS FOR SPE PURIFICATION

4.1 <u>Materials and equipment required</u>

In addition to standard laboratory equipment, the following material is required for the use of **AFFINIMIP® SPE** cartridges:

- SPE vacuum manifold or a SPE automate
- Nitrogen Mini-vap evaporator or Speed vac to dry the collected samples



- Distilled or deionized water
- Methanol (MeOH)
- Acetic acid (AA)
- Phosphate buffer solution pH 7 (I = 10mM) or (Phosphoric acid 1mol/L (H₃PO₄) and Sodium hydroxide 1mol/L (NaOH))

4.2 Flow rate

It is very important to follow the flow rate given in the protocol.

Most especially for the loading, if the sample flow rate is too high, components may not interact sufficiently with the sorbent and the analyte recovery yield will be lower.

4.3 <u>Drying step</u>

If the drying step is not correctly applied, a decreasing of recoveries is observed.

4.4 <u>Template use</u>

For the preparation of the MIP, a template is required. It has a molecular weight of 225 g/mol.

5 PURIFICATION PROCEDURE OF METANEPHRINES FROM PLASMA OR SERUM

5.1 <u>Preparation of solutions</u>

- Solution (60/40) Water/Methanol (v/v)

In a 10mL-volumetric flask, add 4mL of Methanol and complete with water.

- Phosphate Buffer pH=7 (I=10mM)

In a 100mL-volumetric flask, mix 510 μ L of H₃PO₄ (1mol/L), 760 μ L NaOH (1mol/L) in Water. Adjust to pH 7.

- Methanol with 5% Acetic acid

In a 100mL-volumetric flask, add 5mL of acetic acid and complete with Methanol.

5.2 <u>Preparation of the loading solution from plasma or serum (dilution by 5)</u>

In a 5mL volumetric flask, add 1mL of plasma or serum and complete with water.

This solution is used as the loading solution.



5.3 <u>Protocol for purification of Metanephrines</u>

Step (Flow rate)	AFFINIMIP® SPE Metanephrines (30mg/1mL)
Equilibration with (2 drops/s)	 1 mL phosphate buffer pH 7 2 mL deionized water Do not allow the cartridge to dry during conditioning
Loading (L) (1 drop every 2 seconds)	500μL to 1500μL of loading solution
Washing of interferents (W1) (1 drop/s)	1mL Water 500µL (60/40) Water/Methanol
Drying	Apply vacuum through cartridge during 30 seconds
Washing of interferents (W2) (1 drop/s)	500µL Methanol
Elution (E) (1drop/s)	1mL Methanol / Acetic acid (5%)

The extracts (E) are evaporated until dryness under nitrogen with a mini-vap evaporator at room temperature (or a centrifugal evaporator). These residues are dissolved in mobile phase for further analysis. Alternatively, the elution may be diluted to a known volume by addition of water for further analysis.



PRODUCTS LIST

AFFINIMIP® SPE Products	Designation	Description
Multimyco10	AFFINIMIP® SPE Multimyco10	selective SPE cartridges 3mL for ZON, OTA, HT-2, T-2, Aflatoxins and Fumonisins
Zearalenone	AFFINIMIP® SPE Zearalenone	selective SPE cartridges 3mL for ZON
Ochratoxin A	AFFINIMIP® SPE Ochratoxin A	selective SPE cartridges 3mL for OTA
	AFFINIMIP® SPE Patulin	selective SPE cartridges for Patulin
Patulin	AFFINIMIP® SPE Patulin & Pectinase kit	kit of selective SPE cartridges for Patulin + 50mL pectinase enzyme solution
Deoxynivalenol	AFFINIMIP® SPE Deoxynivalenol	selective SPE cartridges 6mL for DON
Phenolics	AFFINIMIP® SPE Phenolics	selective SPE cartridges for Phenolic compounds
Estrogens	AFFINIMIP® SPE Estrogens	selective SPE cartridges for Estrogens
Zeranol Residues	AFFINIMIP® SPE Zeranol Residues	selective SPE cartridges for Zeranol Residues
Bisphenol A	AFFINIMIP® SPE Bisphenol A	Bisphenol A
FumoZON	AFFINIMIP® SPE FumoZON	selective SPE cartridges for Fumonisins and Zearalenone
Chloramphenicol	AFFINIMIP® SPE Chloramphenicol	selective SPE cartridges for Chloramphenicol
Tamoxifen	AFFINIMIP® SPE Tamoxifen	selective SPE cartridges for Tamoxifen
Catecholamines	AFFINIMIP® SPE Catecholamines AFFINIMIP® SPE	selective SPE cartridges for Catecholamines
Metanephrines	Catecholamines AFFINIMIP® SPE	selective SPE cartridges for Catecholamines selective SPE cartridges for Metanephrines
Amphetamines	Metanephrines AFFINIMIP® SPE	selective SPE cartridges for Amphetamines
PECTINASE	Amphetamines Pectinase solution	50 mL pectinase enzyme solution
AttractSPE™ Products	Designation	Description
w/o	AttractSPE™ W/O	HLB SPE cartridges sorbent
SCX	AttractSPE™ SCX	Strong Cation Exchange SPE cartridges sorbent
wcx	AttractSPE™ WCX	Weak Cation Exchange SPE cartridges sorbent
SAX	AttractSPE™ SAX	Strong Anion Exchange SPE cartridges sorbent
WAX	AttractSPE™ WAX	Weak Anion Exchange SPE cartridges sorbent
DVB	AttractSPE™ <mark>DVB</mark>	Reversed Phase Copolymer SPE cartridges sorbent
Anionic & Cationic AttractSPE polymeric cartridges	AttractSPE™ KIT	Kit of 10 cartridges of each sorbent (SAX, WAX, WCX, SCX)

For more information:

For more information on our products & services, please visit www.polyintell.com